**Bold type** indicates authors of commentaries, editorials, and Resource Corners; (No.) indicates issue number.


Abdelnoor M, 34
**Agarwal JB, 31**
Agnelli M, 21
**Albanese A, 14**
Allan JC, 41
Annane D, 44
Antithrombotics in the Secondary Prevention of Events in Coronary Thrombosis-2 Research Group, 9
Arnesen H, 34
**Atkins D, 6-7**
Avenell A, 59
Avierinos J-F, 76
Avin M, 13

Badamgarav E, 55
Bailin SJ, 62
Baines CJ, 38
Band PA, 20
**Barbano R, 47**
Barbui C, 52
Barrett-Connor, 7
Bausell RB, 21
**Beigel J, 44**
Bent-Hansen J, 53
Berman BM, 21
Bermingham JM, 24
Bernard B, 15
BiCard Investigators, 8
**Bijlsma J, 19**
Bittner V, 6
Bjorkman D, 50
Bombardier C, 51
Bonnefoy E, 31
**Bounameaux H, 24**
Brandt LJ, 50
Brashear A, 22
Breathing Not Properly Multinational Study Investigators, 23
Broadley KE, 72
**Bronson DL, 12**
**Brott BC, 76**
Brousseau L, 43
Brown CD, 73
Brown MD, 24
Bryson CL, 80
**Bucher HC, A-11-12 (No. 2)**
**Burdette SD, A-9-10 (No. 1)**
Burgio KL, 75
Buschke H, 25

Cadranel JF, 15
Camargo CA Jr, 18
Carballo D, 83
Chan BK, 39, 40
Chan KJ, 67
Charpentier C, 44
Cheang M, 69
**Cheung BM, 2-3, A-15 (No. 2)**
Chiba N, 17
Chu KM,11
**Cina CS, 66**
Clarke R, 79
Clarkson JE, 46
Coffey CS,4
**Cohn JN, 64**
**Colditz G, 40-41**
Connolly SJ, 36
Crowther MA, 10
Crystal E, 36

Dahlöf B, 37
Davies AN, 72
Davis JC Jr, 19
De Feyter P, 1
de Gans J, 60
De Vet HC, 47
Deckers JW, 9
**Delaney B, 16-17**
**Denman AM, 77**
Devereaux RB, 37
Dickstein K, 65
Doig CJ, 81
Donaldson C, 81
Dose Adjustment for Normal Eating, 68
**Douketis JD, 10, 35**
Dubnov G, 63
Ducharme FM, 42

Eberhardt K, 77
Eckardt L, 61
Eden OB, 46
Edmonds ML, 18
Eikelboom BC, 54
**Eikelboom J, 33, 79**
Elovic E, 22
Erisdden J, 34
Etminan M, 12
**Evans AT, 68**

Fennerty MB, 50
Fink HA, 5
Flammer Y, 33
Fox KA, 32
Furberg C, 7
Furukawa TA, 52

Geborek P, 77
Geibel A, 35
Gerritsen AA, 47
Gersh BJ, 76
Gibbs JS, 72
**Gibson PG, 43**
Gillen D, 16
**Gillespie WJ, 49**
Ginger T, J36
Goetz C, 14
Goldhaber SZ, 24
Goode PS, 75
Gordon MF, 22
Gorman JD, 19
Goulenok C, 15
Grady D, 6
Grobbee DE, 9
Grozinsky S, 45

Haffajee CI, 62
**Hankey GJ, 22**
Hansen HL, 53
Hanson J, 71
Harber P, 48
Harding GK, 69
**Harrington RA, 1**
**Havranek E, 8**
Hawkey CJ, 51
**Haynes RB, A-11 (No. 1)**
He YD, 82
**Hebert PR, 4, 30**
Heinrich R, 35
Helfand M, 39
Henderson RA, 32
Henning JM, 55
**Henry D, 51**
Herrington D, 6
HERS Research Group, 6-7
Hess OM, 33
Heusel G, 35
Hicks GC, 42
Higgins JP, 72
**Hillegass WB, 32**
**Hillner BE, 82**
**Hoffman RA, 74**
Homocysteine Studies Collaboration, 78
Hopkins RO, 67
**Hotopf M, 53**
**Howden CW, 11**
Howe RW, 74
**Hudson MP, 9**
Hulley S, 7
Humphrey LL, 39, 40-41
Hurlen M, 34
Hurwitz EL, 48

Ioannou GN, 80

**Jain M, 63**
Jennings AL, 72
Johnson D, 81
Jonker JJ, 9
Junod AF, 83

Kasper W, 35
Katz M, 25
**Kazlauskaite R, 68**
Kerber RE, 8
Kirchhof P, 61
Kjekshus J, 65
Kjeldsen SE, 37
Klerk M, 79
Klysner R, 53

**Ko DT, 4, 30**
**Kolias TJ, 83**
Koller W, 14
Konstantinides S, 35
Krishnaswamy P, 23
**Kronzon I, 36**
**Kudva YC, 69**
**Kumana CR, 2-3, A-15 (No. 2)**
Kuslansky G, 25

Lacasse Y, 43
**Lader E, 23**
Lai KC, 11
Laine L, 51
Lam SK, 11
Lapostelle F, 31
**Lauder IJ, 2-3, A-15 (No. 2)**
**Laupacis A, 10**
Lee H, 81
**Lee HN, A-13 (No. 2)**
Lee W-L, 21
Leibovici L, 45
Leizorovicz A, 31
Lescol Intervention Prevention Study, 1
Levi Z, 13
Levine MA, 12
Lindqvist E, 77
**Lip GYH, 62**
Lipton RB, 25
Locher JL, 75
**Logan RF, 80**
Loh P, 61

Macaya C, 1
MacDonald R, 5, 74
Maisel AS, 23
Mali WP, 54
Manns BJ, 81
**Mark A, A-11-12 (No. 2)**
McColl KE, 16
McCrory DC, 73
**McDonald JWD, 71**
**McGettigan P, 51**
McGuire H, 52
Melton LJ 3rd, 76
MRC/BHF Heart Protection Study, 2-3
Miller AB, 38
**Mills CS, 55**
Milne AC, 59
Milne S, 43
**Mitchell AM, 59**
**Molloy DW, 25**
Morgenstern H, 48
Moseley JB, 49
**Mulrow CD, 65**
Multicentre Aneurysm Screening Study Group, 66
Murray LS, 16

Nederkoorn PJ, 54
Nelson DF, 5
Nelson HD, 41
**Newman D, 61**
Niaz MA, 63
**Nicolle L, 46,** 69
**Niewoehner DW, 73**
Nowak RM, 23
Nygren P, 41

O'Malley, 49
OPTIMAAL Steering Committee, for the OPTIMAAL Study Group, 65

Page RL, 8
Paul M, 45
**Peters J, 18**
Petersen NJ, 49
Pfeffer M, 64
**Phillips P, 13**
Pin R, 33
**Pitts S, 67**
Poewe W, 14
Pollack CV Jr, 18
Poole-Wilson PA, 32
Potter J, 59
Pruitt R, 71

Rabeneck L, 70
Rachmani R, 13
**Rambaldini G, A-12 (No. 1)**
**Rao G, A-11-12 (No. 3)**
Rascol O, 14
Ravid M, 13
**Ray JG, 78**
Reeves MJ, 24
**Reichen J, 15**
**Rembold CM, 37**
Reynauld JP, 20
**Richardson WS, A-9-10 (No. 1)**
**Robbins RA, 72**
Rochon PA, 12
Rockey DC, 80
Roffi M, 33
**Rosenberg M, 63**
**Rotstein C, 45**
Rowe BH,18, 24
**Rubenstein JH, 50**
Russell JK, 8
Rutks IR, 5, 74

Sébille V, 44
Sack KE, 19
Safdi M, 71
Sampaio C, 14
**Sander O, 21**
Sarasin FP, 83
Saxne T, 77

Schnurr T, 10
Schnyder G, 33
Scholten RJ, 47
Serruys PW, 1
**Shafran SD, 60**
**Shekelle P, 48**
**Shufelt K, A-14 (No. 2)**
**Simon G, 52**
Sinclair P, 17
Singh RB, 63
Slavachevski I, 13
Sleik K, 36
Sliwinksi M, 25
Smith P, 34
Soares-Weiser K, 45
Soeken KL, 21
Somberg JC, 62
Souchek J, 70
Sox HC, 40
**Straus SE, A-13 (No. 1)**
**Swann AC, 4, 30**
**Symmons D, 20**

**Talley NJ, 70**
Teo KK, 64
Teutsch SM, 41
**Thacker HL, 75**
**Thien F, 42**
To T, 38
Tomlinson G, 12
**Topol EJ, 34**
Torrance GW, 20

van de Beek D, 60
van de Vijver MJ, 82
van Es RF, 9
van´t Veer LJ, 82
Veldhuyzen van Zanten SJ, 17
Verheugt FW, 9
Verhoef P, 79

Wall C, 38
**Wardlaw JM, 54**
Weaver LK, 67
Weingarten SR, 55
Weiss NS, 80
**Wiedemann HP, 81**
**Willett LR, 38-39**
**Wilson SA, 5**
Wilt TJ, 5, 74
Woolf SH, 39
Worthington HV, 46
Wristers K, 70

Yusef S, 36, 64

Zhanel GG, 69

A

*ACP Journal Club*
ACP Journal Club sharpens the focus on studies of interest to internists (Editorial), A-11 (No. 1)
Adrenal cortex hormones
Review: Inhaled corticosteroids reduce hospital admissions in acute asthma, 18
Adrenal gland hypofunction
Hydrocortisone and fludrocortisone improved 28-day survival in septic shock and adrenal insufficiency, 44
Adrenergic alpha-antagonists
Review: Terazosin improves urologic symptoms in benign prostatic hyperplasia, 74
Adrenergic beta-antagonists
Review: β-blockers increase fatigue and sexual dysfunction but not depression after myocardial infarction, 4
Review: β-blockers increase fatigue and sexual dysfunction but not depression after myocardial infarction (Letter), 30
Review: β-blockers, sotalol, amiodarone, and biatrial pacing all reduce atrial fibrillation after heart surgery, 36
Alzheimer disease
The Memory Impairment Screen more accurately screened for Alzheimer disease than the 3-word memory test in older adults, 25
Aminosalicylic acids
Balsalazide achieved symptomatic remission sooner than mesalamine for ulcerative colitis, 71
Amiodarone
Intravenous water-soluble amiodarone improved 24-hour survival in incessant ventricular tachycardia, 62
Review: β-blockers, sotalol, amiodarone, and biatrial pacing all reduce atrial fibrillation after heart surgery, 36
Anemia, iron-deficiency
Men and postmenopausal women with iron deficiency had increased risk for gastrointestinal malignancy, 80
Angina pectoris
An Indo-Mediterranean diet was more effective than a control diet in primary and secondary coronary artery disease prevention, 63
An interventional strategy was better than a conservative strategy in unstable angina or non–ST-elevation MI, 32
Angioplasty, transluminal, percutaneous coronary
Fluvastatin reduced major adverse cardiac outcomes after first percutaneous coronary intervention, 1
Homocysteine-lowering therapy improved outcomes after percutaneous coronary intervention, 33
Prehospital fibrinolysis was as good as primary angioplasty after myocardial infarction, 31
Angiotensin-converting enzyme inhibitors
Review: Concomitant aspirin use does not reduce the effectiveness of ACE inhibitors, 64
Anti-arrhythmia agents
Intravenous water-soluble amiodarone improved 24-hour survival in incessant ventricular tachycardia, 62
Antiasthmatic agents
Review: Inhaled corticosteroids reduce hospital admissions in acute asthma, 18
Anticoagulants
Coumadin alone or aspirin plus coumadin reduced coronary events and death after acute myocardial infarction or unstable angina, 9
Warfarin alone or with aspirin was superior to aspirin alone after acute myocardial infarction but increased bleeding, 34
Antidepressive agents, tricyclic
Review: Low dose was as effective as standard dose tricyclic antidepressants in adults with depression, 52
Antifungal agents
Review: Antifungals absorbed or partially absorbed from the GI tract prevent oral candidiasis in cancer patients receiving treatment, 46
Antihypertensive agents
Losartan was more effective than atenolol for isolated systolic hypertension with left ventricular hypertrophy, 37
Anti-inflammatory agents, non-steroidal
Review: Antileukotrienes are less effective than inhaled corticosteroids in chronic asthma, 42
Review: Inhaled corticosteroids reduce hospital admissions in acute asthma, 18
Review: S-adenosylmethionine treats osteoarthritis as effectively as nonsteroidal anti-inflammatory drugs, 21
Upper gastrointestinal event risk with cox-2 inhibitors depended on known risk factors, 51
Antioxidants
Antioxidant vitamins did not reduce death, vascular events, or cancer in high-risk patients, 3
Antirheumatic agents
Etanercept was effective and safe for ankylosing spondylitis, 19
Long-term prognosis, measured by health status, disease process, and damage, varies in early rheumatoid arthritis, 77
Aortic aneurysm, abdominal
Screening for abdominal aortic aneurysm reduced death from AAA in older men, 66
Arthritis, rheumatoid
Long-term prognosis, measured by health status, disease process, and damage, varies in early rheumatoid arthritis, 77
Upper gastrointestinal event risk with COX-2 inhibitors depended on known risk factors, 51
Arthroscopy
Arthroscopic surgery was not effective for relieving pain or improving function in osteoarthritis of the knee, 49
Aspirin
Coumadin alone or aspirin plus coumadin reduced coronary events and death after acute myocardial infarction or unstable angina, 9
Lansoprazole reduced recurrence of ulcer complications in long-term use of low-dose aspirin, 11
Review: Concomitant aspirin use does not reduce the effectiveness of ACE inhibitors, 64
Warfarin alone or with aspirin was superior to aspirin alone after acute myocardial infarction but increased bleeding, 34
Asthma
Review: Antileukotrienes are less effective than inhaled corticosteroids in chronic asthma, 42
Review: Inhaled corticosteroids reduce hospital admissions in acute asthma, 18
Atenolol
Losartan was more effective than atenolol for isolated systolic hypertension with left ventricular hypertrophy, 37
Atrial fibrillation
Anterior–posterior electrode position led to more cardioversion success in atrial fibrillation, 61

Biphasic waveform shocks were effective and efficient for cardioversion of atrial fibrillation, 8
Review: β-blockers, sotalol, amiodarone, and biatrial pacing all reduce atrial fibrillation after heart surgery, 36
Atrial natriuretic factor
BNP levels had high sensitivity but moderate specificity for detecting congestive heart failure in the emergency department, 23

B

Bacteriuria
Antimicrobial treatment did not reduce complications of urinary tract infection in diabetes and asymptomatic bacteriuria, 69
Behavior therapy
Self-administered behavioral training was equal to training plus biofeedback or verbal feedback in urge incontinence, 75
Biofeedback
Self-administered behavioral training was equal to training plus biofeedback or verbal feedback in urge incontinence, 75
Blood coagulation disorders
Oral vitamin K lowered reversed warfarin-associated coagulopathy faster than subcutaneous vitamin K, 10
Botulinum toxin type A
Botulinum toxin A injections improved wrist and finger spasticity after stroke, 22
Breast neoplasms
Annual screening with mammography and breast examination did not reduce breast cancer mortality in women 40 to 49 years of age, 38
A gene-expression profile independently predicted outcome of disease in young women with breast cancer, 82
Review: Mammography reduces breast cancer mortality rates, 39
Breath tests
Noninvasive Helicobacter pylori testing was as effective as endoscopy for managing dyspepsia, 16
Bronchodilator agents
Review: Ipratropium is not more effective than β-agonists for acute exacerbations of chronic obstructive pulmonary disease, 73

C

Cancer
Antioxidant vitamins did not reduce death, vascular events, or cancer in high-risk patients, 3
Candidiasis, oral
Review: Antifungals absorbed or partially absorbed from the GI tract prevent oral candidiasis in cancer patients receiving treatment, 46
Captopril
Losartan did not differ from captopril for reducing all-cause mortality after acute myocardial infarction, 65
Carbon monoxide poisioning
3 hyperbaric oxygen treatments reduced cognitive sequelae of acute carbon monoxide poisoning, 67
Cardiotonic agents
An interventional strategy was better than a conservative strategy in unstable angina or non–ST-elevation MI, 32
Cardiovascular disease
Antioxidant vitamins did not reduce death, vascular events, or cancer in high-risk patients, 3
Long-term mortality and cardiovascular morbidity increased in some patients with asymptomatic mitral valve prolapse, 76
MRC/BHF Heart Protection Study (Letter), A-13 (No. 2)
Review: Elevated homocysteine levels are modestly associated with increased ischemic heart disease and stroke risk, 78
Simvastatin reduced mortality and vascular events in high-risk patients, 2
Carotid stenosis
The combined use of duplex ultrasonography and magnetic resonance angiography most accurately diagnosed carotid artery stenosis, 54
Carpal tunnel syndrome
Surgery was associated with greater long-term treatment success than wrist splinting in the carpal tunnel syndrome, 47
Cerebrovascular accident
Review: Elevated homocysteine levels are modestly associated with increased ischemic heart disease and stroke risk, 78
Chiropractic
Cervical spine manipulation was not better than mobilization for improving outcomes in neck pain, 48
Cholinergic antagonists
Review: Ipratropium is not more effective than $\beta_2$-agonists for acute exacerbations of chronic obstructive pulmonary disease, 73
Chronic disease
Review: Most disease management programs for providers and patients lead to improvements in care, 55
Cirrhosis
Review: Flumazenil leads to clinical and electroencephalographic improvement in hepatic encephalopathy in patients with cirrhosis, 15
Citalopram
Prophylactic citalopram reduced recurrences of unipolar major depression, 53
Cognition disorders
3 hyperbaric oxygen treatments reduced cognitive sequelae of acute carbon monoxide poisoning, 67
Colitis, ulcerative
Balsalazide achieved symptomatic remission sooner than mesalamine for ulcerative colitis, 71
Colonic diseases, functional
Review: Tegaserod or alosetron is effective for the irritable bowel syndrome, 50
Computers, handheld
Practice Corner: Clinical practice guidelines and handheld computers, A-11 (No. 3)
Practice Corner: Taking evidence in hand, A-9 (No. 1)
Coronary disease
Estrogen plus progestin was not effective for long-term secondary prevention of coronary heart disease in postmenopausal women, 6
*Evidence-Based Cardiology* (Resource Corner), A-16 (No. 2)
An interventional strategy was better than a conservative strategy in unstable angina or non–ST-elevation MI, 32
Review: MTHFR TT genotype increases risk for coronary heart disease, 79
Cost–benefit analysis
Activated protein C was cost-effective for prolonging survival in a subgroup of patients with severe sepsis, 81
Hylan G-F 20 was safe and effective in knee osteoarthritis and had a relatively low cost–utility ratio, 20
Cystitis
Antimicrobial treatment did not reduce complications of urinary tract infection in diabetes and asymptomatic bacteriuria, 69

## D

Decompression, surgical
Surgery was associated with greater long-term treatment success than wrist splinting in the carpal tunnel syndrome, 47
Depressive disorder
Prophylactic citalopram reduced recurrences of unipolar major depression, 53
Review: Low dose was as effective as standard dose tricyclic antidepressants in adults with depression, 52
Dexamethasone
Dexamethasone improved disability in acute bacterial meningitis, 60
Diabetes mellitus
Antimicrobial treatment did not reduce complications of urinary tract infection in diabetes and asymptomatic bacteriuria, 69
Involving patients with type 2 diabetes, hypertension, and hyperlipidemia in disease management improved outcomes, 13
MRC/BHF Heart Protection Study (Letter), A-13 (No. 2)
Simvastatin reduced mortality and vascular events in high-risk patients, 2
Training in flexible intensive insulin management improved glycemic control and quality of life in type 1 diabetes, 68
Diabetic angiopathies and nephropathies
Involving patients with type 2 diabetes, hypertension, and hyperlipidemia in disease management improved outcomes, 13
Diet
Indo-Mediterranean diet was more effective than a control diet in primary and secondary coronary artery disease prevention, 63
Training in flexible intensive insulin management improved glycemic control and quality of life in type 1 diabetes, 68
Dietary proteins
Review: Oral protein and energy supplements reduce all-cause mortality in elderly persons, 59
Dietary supplements
Review: S-adenosylmethionine treats osteoarthritis as effectively as nonsteroidal anti-inflammatory drugs, 21
Disability evaluation
Long-term prognosis, measured by health status, disease process, and damage, varies in early rheumatoid arthritis, 77
Disease-free survival
Long-term mortality and cardiovascular morbidity increased in some patients with asymptomatic mitral valve prolapse, 76
Dyspepsia
*Helicobacter pylori* eradication improved dyspepsia symptoms, 17
Noninvasive *Helicobacter pylori* testing was as effective as endoscopy for managing dyspepsia, 16
Omeprazole was effective in the short term for uninvestigated dyspepsia, 70
Dyspnea
BNP levels had high sensitivity but moderate specificity for detecting congestive heart failure in the emergency department, 23
Review: Oral or parenteral opioids alleviate dyspnea in palliative care, 72

## E

Echocardiography
Echocardiography was useful in evaluating unexplained syncope, 83
Electric countershock
Anterior–posterior electrode position led to more cardioversion success in atrial fibrillation, 61
Biphasic waveform shocks were effective and efficient for cardioversion of atrial fibrillation, 8
Electrodes
Anterior–posterior electrode position led to more cardioversion success in atrial fibrillation, 61
Emergency medical services
Prehospital fibrinolysis was as good as primary angioplasty after myocardial infarction, 31
Energy intake
Review: Oral protein and energy supplements reduce all-cause mortality in elderly persons, 59
Enzyme-linked immunosorbent assay
Review: ELISA D-dimer is sensitive but not specific in diagnosing pulmonary embolism in an ambulatory clinical setting, 24
Estrogen replacement therapy
Estrogen plus progestin was not effective for long-term secondary prevention of coronary heart disease in postmenopausal women, 6
Long-term estrogen-plus-progestin increased venous thromboembolism and biliary tract surgery in postmenopausal women, 7
Evidence-based medicine
*Clinical Evidence Concise* (Resource Corner), A-12 (No. 1)
*Evidence-Based Cardiology* (Resource Corner), A-16 (No. 2)
*Evidence-Based Herbal Medicine* (Resource Corner), A-13 (No. 1)
National Electronic Library of Health (Resource Corner), A-15 (No. 2)
Numbers needed to treat derived from meta-analysis: a word of caution (Editorial), A-11 (No. 2)
Practice Corner: Clinical practice guidelines and handheld computers, A-11 (No. 3)
Practice Corner: Taking evidence in hand (Editorial), A-9 (No. 1)
Exercise tolerance
Review: Respiratory rehabilitation improves health-related quality of life in chronic obstructive pulmonary disease, 43

## F

Feedback
Review: Most disease management programs for providers and patients lead to improvements in care, 55
Fibrinolytic agents
Heparin plus alteplase reduced morbidity more than heparin alone in submassive pulmonary embolism, 35
Fludrocortisone
Hydrocortisone and fludrocortisone improved 28-day survival in septic shock and adrenal insufficiency, 44
Flumazenil
Review: Flumazenil leads to clinical and electroencephalographic improvement in hepatic encephalopathy in patients with cirrhosis, 15
Folic acid
Homocysteine-lowering therapy improved outcomes after percutaneous coronary intervention, 33

## G

Gastrointestinal diseases
Men and postmenopausal women with iron deficiency had increased risk for gastrointestinal malignancy, 80
Upper gastrointestinal event risk with COX-2 inhibitors depended on known risk factors, 51
Gastroscopy
Noninvasive *Helicobacter pylori* testing was as effective as endoscopy for managing dyspepsia, 16

Gene-expression profiling
A gene-expression profile independently predicted outcome of disease in young women with breast cancer, 82

H

Headache
Review: Angiotensin II receptor antagonists prevent headache in patients with mild-to-moderate hypertension, 12
Health status indicators
Long-term prognosis, measured by health status, disease process, and damage, varies in early rheumatoid arthritis, 77
Heart diseases
Echocardiography was useful in evaluating unexplained syncope, 83
Heart failure, congestive
BNP levels had high sensitivity but moderate specificity for detecting congestive heart failure in the emergency department, 23
Review: β-blockers increase fatigue and sexual dysfunction but not depression after myocardial infarction, 4
Review: β-blockers increase fatigue and sexual dysfunction but not depression after myocardial infarction (Letter), 30
Review: Concomitant aspirin use does not reduce the effectiveness of ACE inhibitors, 64
*Helicobacter pylori*
*Helicobacter pylori* eradication improved dyspepsia symptoms, 17
Lansoprazole reduced recurrence of ulcer complications in long-term use of low-dose aspirin, 11
Noninvasive *Helicobacter pylori* testing was as effective as endoscopy for managing dyspepsia, 16
Heparin
Heparin plus alteplase reduced morbidity more than heparin alone in submassive pulmonary embolism, 35
Hepatic encephalopathy
Review: Flumazenil leads to clinical and electroencephalographic improvement in hepatic encephalopathy in patients with cirrhosis, 15
Homocysteine
Homocysteine-lowering therapy improved outcomes after percutaneous coronary intervention, 33
Review: Elevated homocysteine levels are modestly associated with increased ischemic heart disease and stroke risk, 78
Hormone replacement therapy
Review: Observational studies adjusting for socioeconomic status and lifestyle show no association between HRT and CAD, 40
Review: Risks and benefits of HRT comparing various sources of evidence, 41
Hydrochlorothiazide
Losartan was more effective than atenolol for isolated systolic hypertension with left ventricular hypertrophy, 37
Hydrocortisone
Hydrocortisone and fludrocortisone improved 28-day survival in septic shock and adrenal insufficiency, 44
Hydroxymethylglutaryl-CoA reductase inhibitors
Fluvastatin reduced major adverse cardiac outcomes after first percutaneous coronary intervention, 1
Hyperbaric oxygenation
3 hyperbaric oxygen treatments reduced cognitive sequelae of acute carbon monoxide poisoning, 67
Hyperlipidemia
Involving patients with type 2 diabetes, hypertension, and hyperlipidemia in disease management improved outcomes, 13
Hypertension
Involving patients with type 2 diabetes, hypertension, and hyperlipidemia in disease management improved outcomes, 13
Losartan was more effective than atenolol for isolated systolic hypertension with left ventricular hypertrophy, 37
Review: β-blockers increase fatigue and sexual dysfunction but not depression after myocardial infarction, 4
Review: β-blockers increase fatigue and sexual dysfunction but not depression after myocardial infarction (Letter), 30

I

Impotence
Review: Sildenafil improves erectile dysfunction and is well tolerated, 5
Insulin
Training in flexible intensive insulin management improved glycemic control and quality of life in type 1 diabetes, 68
International normalized ratio
Oral vitamin K lowered reversed warfarin-associated coagulopathy faster than subcutaneous vitamin K, 10
Internet
National Electronic Library of Health (Resource Corner), A-15 (No. 2)
Practice Corner: Taking evidence in hand, A-9 (No. 1)
Intestinal absorption
Review: Antifungals absorbed or partially absorbed from the GI tract prevent oral candidiasis in cancer patients receiving treatment, 46
Ipratropium
Review: Ipratropium is not more effective than $\beta_2$-agonists for acute exacerbations of chronic obstructive pulmonary disease, 73
Iron
Men and postmenopausal women with iron deficiency had increased risk for gastrointestinal malignancy, 80

L

Lactams
Review: β-lactam alone was better than β-lactam plus aminoglycoside for cancer patients with neutropenia, 45
Leukotriene antagonists
Review: Antileukotrienes are less effective than inhaled corticosteroids in chronic asthma, 42
Lidocaine
Intravenous water-soluble amiodarone improved 24-hour survival in incessant ventricular tachycardia, 62
Losartan
Losartan did not differ from captopril for reducing all-cause mortality after acute myocardial infarction, 65
Losartan was more effective than atenolol for isolated systolic hypertension with left ventricular hypertrophy, 37

M

Magnetic resonance angiography
The combined use of duplex ultrasonography and magnetic resonance angiography most accurately diagnosed carotid artery stenosis, 54
Mammography
Annual screening with mammography and breast examination did not reduce breast cancer mortality in women 40 to 49 years of age, 38
Review: Mammography reduces breast cancer mortality rates, 39

Manipulation, chiropractic
Cervical spine manipulation was not better than mobilization for improving outcomes in neck pain, 48
Mass screening
Annual screening with mammography and breast examination did not reduce breast cancer mortality in women 40 to 49 years of age, 38
Review: Mammography reduces breast cancer mortality rates, 39
Screening for abdominal aortic aneurysm reduced death from AAA in older men, 66
Medicine, herbal
*Evidence-Based Herbal Medicine* (Resource Corner), A-13 (No. 1)
Memory
The Memory Impairment Screen more accurately screened for Alzheimer disease than the 3-word memory test in older adults, 25
Meningitis, bacterial
Dexamethasone improved disability in acute bacterial meningitis, 60
Mesalamine
Balsalazide achieved symptomatic remission sooner than mesalamine for ulcerative colitis, 71
Meta-analysis
Numbers needed to treat derived from meta-analysis: a word of caution (Editorial), A-11 (No. 2)
Migraine
Practice Corner: Clinical practice guidelines and handheld computers, A-11 (No. 3)
Mitral valve prolapse
Long-term mortality and cardiovascular morbidity increased in some patients with asymptomatic mitral valve prolapse, 76
Muscle spasticity
Botulinum toxin A injections improved wrist and finger spasticity after stroke, 22
Myocardial infarction
Coumadin alone or aspirin plus coumadin reduced coronary events and death after acute myocardial infarction or unstable angina, 9
Estrogen plus progestin was not effective for long-term secondary prevention of coronary heart disease in postmenopausal women, 6
An Indo-Mediterranean diet was more effective than a control diet in primary and secondary coronary artery disease prevention, 63
An interventional strategy was better than a conservative strategy in unstable angina or non–ST-elevation MI, 32
Long-term estrogen-plus-progestin increased venous thromboembolism and biliary tract surgery in postmenopausal women, 7
Losartan did not differ from captopril for reducing all-cause mortality after acute myocardial infarction, 65
Prehospital fibrinolysis was as good as primary angioplasty after myocardial infarction, 31
Review: β-blockers increase fatigue and sexual dysfunction but not depression after myocardial infarction, 4
Review: β-blockers increase fatigue and sexual dysfunction but not depression after myocardial infarction (Letter), 30
Warfarin alone or with aspirin was superior to aspirin alone after acute myocardial infarction but increased bleeding, 34
Myocardial ischemia
Fluvastatin reduced major adverse cardiac outcomes after first percutaneous coronary intervention, 1

N

Narcotics
Review: Oral or parenteral opioids alleviate dyspnea in palliative care, 72
Neck pain
Cervical spine manipulation was not better than mobilization for improving outcomes in neck pain, 48
Neoplasms
A gene-expression profile independently predicted outcome of disease in young women with breast cancer, 82
Review: β-lactam alone was better than β-lactam plus aminoglycoside for cancer patients with neutropenia, 45
Neuropsychological tests
The Memory Impairment Screen more accurately screened for Alzheimer disease than the 3-word memory test in older adults, 25
Neutropenia
Review: β-lactam alone was better than β-lactam plus aminoglycoside for cancer patients with neutropenia, 45
Nutritional disorders
Review: Oral protein and energy supplements reduce all-cause mortality in elderly persons, 59

O

Omeprazole
Lansoprazole reduced recurrence of ulcer complications in long-term use of low-dose aspirin, 11
Omeprazole was effective in the short term for uninvestigated dyspepsia, 70
Osteoarthritis
Arthroscopic surgery was not effective for relieving pain or improving function in osteoarthritis of the knee, 49
Hylan G-F 20 was safe and effective in knee osteoarthritis and had a relatively low cost–utility ratio, 20
Review: S-adenosylmethionine treats osteoarthritis as effectively as nonsteroidal anti-inflammatory drugs, 21
Oxygen inhalation therapy
3 hyperbaric oxygen treatments reduced cognitive sequelae of acute carbon monoxide poisoning, 67

P

Parkinson disease
Review: Several drugs are efficacious for symptomatic treatment of Parkinson disease, 14
Patient compliance
Review: Low dose was as effective as standard dose tricyclic antidepressants in adults with depression, 52
Patient education
Review: Most disease management programs for providers and patients lead to improvements in care, 55
Training in flexible intensive insulin management improved glycemic control and quality of life in type 1 diabetes, 68
Penile erection
Review: Sildenafil improves erectile dysfunction and is well tolerated, 5
Peptic ulcer
Lansoprazole reduced recurrence of ulcer complications in long-term use of low-dose aspirin, 11
Phosphodiesterase inhibitors
Review: Sildenafil improves erectile dysfunction and is well tolerated, 5
Piperazines
Review: Sildenafil improves erectile dysfunction and is well tolerated, 5

Polymorphism
Review: MTHFR TT genotype increases risk for coronary heart disease, 79
Postmenopause
Estrogen plus progestin was not effective for long-term secondary prevention of coronary heart disease in postmenopausal women, 6
Long-term estrogen-plus-progestin increased venous thromboembolism and biliary tract surgery in postmenopausal women, 7
Review: Observational studies adjusting for socioeconomic status and lifestyle show no association between HRT and CAD, 40
Review: Risks and benefits of HRT comparing various sources of evidence, 41
Primary prevention
Review: Observational studies adjusting for socioeconomic status and lifestyle show no association between HRT and CAD, 40
Prostatic hyperplasia
Review: Terazosin improves urologic symptoms in benign prostatic hyperplasia, 74
Protein C
Activated protein C was cost-effective for prolonging survival in a subgroup of patients with severe sepsis, 81
Proton pumps
Omeprazole was effective in the short term for uninvestigated dyspepsia, 70
Pulmonary disease, chronic obstructive
Review: Ipratropium is not more effective than $\beta_2$-agonists for acute exacerbations of chronic obstructive pulmonary disease, 73
Review: Respiratory rehabilitation improves health-related quality of life in chronic obstructive pulmonary disease, 43
Pulmonary embolism
Heparin plus alteplase reduced morbidity more than heparin alone in submassive pulmonary embolism, 35
Review: ELISA D-dimer is sensitive but not specific in diagnosing pulmonary embolism in an ambulatory clinical setting, 24

Q

Quality of life
Hylan G-F 20 was safe and effective in knee osteoarthritis and had a relatively low cost–utility ratio, 20

R

Randomized controlled trials
Numbers needed to treat derived from meta-analysis: A word of caution (Editorial), A-11 (No. 2)
Recall
The Memory Impairment Screen more accurately screened for Alzheimer disease than the 3-word memory test in older adults, 25
Receptors, angiotensin
Review: Angiotensin II receptor antagonists prevent headache in patients with mild-to-moderate hypertension, 12
Review: Risks and benefits of HRT comparing various sources of evidence, 41

S

S-adenosylmethionine
Review: S-adenosylmethionine treats osteoarthritis as effectively as nonsteroidal anti-inflammatory drugs, 21
Sepsis
Activated protein C was cost-effective for prolonging survival in a subgroup of patients with severe sepsis, 81
Serotonin uptake inhibitors
Prophylactic citalopram reduced recurrences of unipolar major depression, 53
Shock, septic
Hydrocortisone and fludrocortisone improved 28-day survival in septic shock and adrenal insufficiency, 44
Simvastatin
MRC/BHF Heart Protection Study (Letter), A-13 (No. 2)
Simvastatin reduced mortality and vascular events in high-risk patients, 2
Sotalol
Review: β-blockers, sotalol, amiodarone, and biatrial pacing all reduce atrial fibrillation after heart surgery, 36
Splints
Surgery was associated with greater long-term treatment success than wrist splinting in the carpal tunnel syndrome, 47
Spondylitis, ankylosing
Etanercept was effective and safe for ankylosing spondylitis, 19
Syncope
Echocardiography was useful in evaluating unexplained syncope, 83

T

Tissue plasminogen activator
Heparin plus alteplase reduced morbidity more than heparin alone in submassive pulmonary embolism, 35
Tumor necrosis factor
Etanercept was effective and safe for ankylosing spondylitis, 19

U

Ultrasonography, Doppler, duplex
The combined use of duplex ultrasonography and magnetic resonance angiography most accurately diagnosed carotid artery stenosis, 54
Urinary incontinence
Self-administered behavioral training was equal to training plus biofeedback or verbal feedback in urge incontinence, 75

V

Vitamin K
Oral vitamin K lowered reversed warfarin-associated coagulopathy faster than subcutaneous vitamin K, 10

W

Warfarin
Oral vitamin K lowered reversed warfarin-associated coagulopathy faster than subcutaneous vitamin K, 10
Warfarin alone or with aspirin was superior to aspirin alone after acute myocardial infarction but increased bleeding, 34